

CERTIFICATE OF EXPRESS MAIL (37 CFR 1.10)

Label No. EE068531687US

I hereby certify that the attached papers and/or fee are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service on the date shown below in an envelope addressed to Box PCT, Assistant Commissioner for Patents, Washington D.C. 20231.

June 9, 2000
Date

Mary Rutkowski
Printed Name

Mary Rutkowski
Signature

FORM PTO-1390 (REV 12/29/99)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

TRANSMITTAL LETTER TO THE UNITED STATES

DESIGNATED/ELECTED OFFICE (DO/EO/US)

CONCERNING A FILING UNDER 35 U.S.C. 371

ATTORNEY'S DOCKET NUMBER

MOG 57701/UST

U. S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/581331

INTERNATIONAL APPLICATION NO.:

PCT/EP98/08162

INTERNATIONAL FILING DATE:

December 10, 1998

PRIORITY DATE CLAIMED:

December 12, 1997

TITLE OF INVENTION:

Constitutive Plant Promoters

EXPRESS MAIL NO.:

EE068531687US

APPLICANT(S) FOR DO/EO/US:

STUIVER, Maarten Hendrik and SIJOLTS, Floor Hendrik

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1)).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371 (c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.

☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☐ Other items or information:

U.S. APPLICATION NO. (if known, see 37 CFR 1.5) <div style="font-size: 24pt; font-weight: bold;">09/581331</div>		INTERNATIONAL APPLICATION NO PCT/EP98/08162		ATTORNEY'S DOCKET NUMBER MOG 57701/UST	
---	--	--	--	---	--

17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492 (a)(1)-(5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$ 970.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$ 840.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO. . . \$ 690.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$ 670.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4). \$ 96.00 <div style="text-align: right;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div>	CALCULATIONS PTO USE ONLY <div style="border: 1px solid black; padding: 2px;"> <div style="display: flex; justify-content: space-between;"> \$ 840.00 </div> </div>
---	--

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	15-20 =	0	x \$18.00	\$ 0	
Independent claims	1- 3 =	0	x \$78.00	\$ 0	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$260.00	\$	
TOTAL OF ABOVE CALCULATIONS =				\$ 840.00	
Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28).				\$	
SUBTOTAL =				\$ 840.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (f)).				+	
TOTAL NATIONAL FEE =				\$ 840.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +				+	
TOTAL FEES ENCLOSED =				\$ 840.00	
				Amount to be refunded:	\$
				Charged	\$

a. ☐ A check in the amount of \$ _____ to cover the above fees is enclosed.


b. ☒ Please charge my Deposit Account No. 09-0013 in the amount of \$ 840.00 to cover the above fees.
A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 09-0013. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

ZENECA Ag Products Inc.
Intellectual Property Section
1800 Concord Pike
P.O. Box 15458
Wilmington, DE 19850-5458



23698

PATENT TRADEMARK OFFICE

Liza D. Hohenschutz
SIGNATURE

June 9, 2000
DATE

Liza D. Hohenschutz
NAME

33,712
REGISTRATION NUMBER

(302)886-1699
TELEPHONE NUMBER

CERTIFICATE OF EXPRESS MAIL (37 CFR 1.10) Label No. EE068531687US

I hereby certify that the attached papers and/or fee are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service on the date shown below in an envelope addressed to: Box Patent Application, Assistant Commissioner for Patents, Washington DC 20231.

June 9, 2000
(Date)

Mary Rutkowski
(Printed Name)

Mary Rutkowski
(Signature)

PATENT APPLICATION
MOG 57701/UST

IN THE UNITED STATES ELECTED OFFICE

INTERNATIONAL APPLICATION NO.: PCT/EP98/08162

INTERNATIONAL FILING DATE: 10 December 1998

PRIORITY DATE CLAIMED: 12 December 1997

APPLICANTS: Maarten Hendrik STUIVER *et al.*

TITLE: CONSTITUTIVE PLANT PROMOTERS

PRELIMINARY AMENDMENT

Box PCT
Assistant Commissioner for Patents
Washington, DC 20231

Sir:

Prior to calculating the filing fee, please amend the above-identified application as follows:

IN THE CLAIMS

3. (Amended) Chimaeric plant promoter according to claim 1 [or 2], characterized in that one promoter of the set of promoters is specifically active in green parts of the plant, while the other promoter is specifically active in underground parts of the plant.

6. (Amended) Chimaeric plant promoter according to claim 4 [or 5], characterized in that the ferredoxin promoter is derived from *Arabidopsis thaliana*.

11. (Amended) Chimaeric plant promoter according to claim 9 [or 10], characterized in that the plastocyanin promoter is derived from *Arabidopsis thaliana*.

12. (Amended) Chimaeric plant promoter according to claim 9[, 10 or 11], characterized in that the S-adenosyl-methionine-1 promoter is derived from *Arabidopsis thaliana*.


15. (Amended) Chimaeric gene construct for the expression of genes in plants comprising the promoter of [any of claims 1-14] claim 1.

REMARKS

The claims have been amended to remove improper multiple dependencies and place them in a better form for examination.

An early and favorable Office Action is requested.

Respectfully submitted,
ZENECA Ag Products



Liza D. Hohenschutz
Reg. No. 33,712
Attorney for Applicants

Dated: June 9, 2000
1800 Concord Pike
P.O. Box 15458
Wilmington, DE 19850-5458
Telephone: (302) 886-1699

g:\intelprop\mog files\mog 57701ust\preliminary amendment.doc

4 ppts

WO 99/31258

09/581331
10 Rec'd PCT/EP98/08162
09 JUN 2000

CONSTITUTIVE PLANT PROMOTERS

FIELD OF THE INVENTION

5 The invention is directed to new plant promoters, more specifically those promoters which can be produced by assembling parts of promoters which have a complementary specificity.

BACKGROUND ART

10 Genetic engineering of plants has become possible by virtue of two discoveries: first of all the possibility of transformation of heterologous genetic material to the plant cell (most efficiently done by the bacterium *Agrobacterium tumefaciens* or related strains) and secondly by the existence of plant promoters which are able to drive
15 the expression of said heterologous genetic material.

A typical plant promoter consists of specific elements. A basis is formed by the minimal promoter element, which enables transcription initiation, often accompanied by a sequence, also denominated as the TATA-box, which serves as a binding place for transcription initiation
20 factors. In most promoters, the presence of this TATA-box is important for proper transcription initiation. It is typically located 35 to 25 basepairs (bp) upstream of the transcription initiation site. Another part of the promoter consists of elements which are able to interact with DNA-binding proteins. Known are G-box binding elements
25 which are based on the hexanucleotide CACGTG motif. These elements have been shown to be able to interact with bZIP DNA-binding proteins which bind as dimers (Johnson & McKnight, Ann. Rev. Biochem, 58, 799-839, 1989). Other G-box related motifs, such as the Iwt and PA motifs have been described (WO 94/12015).

30 These motifs have been shown to be involved in tissue-specific promoter expression in plants. For instance, presence of Iwt tetramers confer embryo-specific expression, while PA tetramers confer high level root expression, low-level leaf expression and no seed expression.

35 Similarly, GT-1 like binding sites (grouped on basis of a moderate consensus sequence GGT^A/A) are described. Such a binding site is found far upstream the promoter region of the *Arabidopsis* plastocyanin promoter and seems to be involved in activation of transcription

during light periods (Fisscher, U. et al., Plant Mol. Biol. 26, 873-886, 1994).

Another sequence-related phenomenon which is found often in plant promoters is the presence of sequences which enable the formation of Z-DNA. Z-DNA is DNA folded in a left-handed helix which is caused by repeats of dinucleotides GC or AC. It is believed that folding in a Z-form influences the availability of the DNA for approach by RNA polymerase molecules, thus inhibiting the transcription rate.

One of the earliest and most important inventions in the field of plant protein expression is the use of (plant) viral and Agrobacterium-derived promoters that provide a powerful and constitutive expression of heterologous genes in transgenic plants. Several of these promoters have been used very intensively in plant genetic research and still are the promoter of choice for rapid, simple and low-risk expression studies. The most famous are the 35S and 19S promoter from Cauliflower Mosaic Virus (CaMV), which was already found to be practically useful in 1984 (EP 0 131 623), the promoters which can be found in the Agrobacterium T-DNA, like the nopaline synthase (nos), mannopine synthase (mas) and octopine synthase (ocs) promoters (EP 0 122 791, EP 0 126 546, EP 0 145 338). A plant-derived promoter with similar characteristics is the ubiquitin promoter (EP 0 342 926).

In time, several attempts have been made to increase the level of expression of these promoters. Examples for this are the double enhanced 35S promoter (US 5,164,316) and, more recently, the superpromoter, which couples parts of the Agrobacterium promoters (EP 729 514).

However, in many cases these promoters do not fulfill the criteria of an ideal promoter. All promoters described above show a clear pattern of organ- or developmental-specific expression, and frequently the pattern of expression found with these promoters is not ideal for some applications. Especially for biotechnological applications like the engineering of fungal and insect resistance, which require expression both in the right location as well as in the right timeframe of plant development there is a need for new

constitutive promoters which are able to give a high level of transgene expression at exactly the right time and place.

SUMMARY OF THE INVENTION

The invention provides for novel plant promoters, characterized in that they comprise 1) a minimal promoter and 2) transcription-activating elements from a set of promoters, which elements direct a complementary pattern and level of transcription in a plant.

More specifically, this plant promoter is a constitutive promoter in which each of the transcription-activating elements do not exhibit an absolute tissue-specificity, but mediate transcriptional activation in most plant parts at a level of $\geq 1\%$ of the level reached in the part of the plant in which transcription is most active. An example of such promoter pairs is a set of promoters in which one is most active in green parts of the plant, while the other promoter is most active in underground parts of the plant. More specifically the new promoter is a combination of the ferredoxin and the *rolD* promoter. Preferably in this construct the minimal promoter element is derived from the ferredoxin promoter and the ferredoxin promoter is derived from *Arabidopsis thaliana*. The *rolD* promoter is derived from *Agrobacterium rhizogenes*.

Also part of the invention is a plant promoter which is a combination of the plastocyanin and the *S*-adenosyl-methionine-1 promoter, whereby preferably the minimal promoter element is derived from the *S*-adenosyl-methionine-1 promoter and both the plastocyanin promoter and the *S*-adenosyl-methionine-1 promoter are derived from *Arabidopsis thaliana*.

Further part of the invention are chimaeric gene constructs for the expression of genes in plants comprising the above disclosed promoters.

DESCRIPTION OF THE FIGURES

Figure 1: Schematic representation of pMOG410 and pMOG1059

Figure 2: Distribution of GUS expression of potato lines transformed with the constructs pMOG1059 en pMOG410. GUS staining was judged visually and classes of expression, relative to the highest GUS expression measured in our lab (set at 4). A value of zero indicates no visible expression.

Figure 3: Graphic representation of the average expression of GUS enzyme in primary transformants of tomato, oilseed rape and potato. GUS expression was determined visually and compared to a high level expressing 35S GUS transgenic tobacco plant ranking a score of 4. Standard deviation of the measured values are indicated on each of the bars.

Figure 4: Graphic representation of the distribution of potato plants with various levels of GUS expression containing SAM-1-, Pc-35S- and PcSAM1-GUS constructs. Scored are expression in leaf mesophyll, leaf vascular system, stem and root.

DETAILED DESCRIPTION OF THE INVENTION

For the purpose of this specification the following definitions are valid:

A **promoter** consists of an RNA polymerase binding site on the DNA, forming a functional transcription initiation start site. A promoter usually consists of at least a TATA box and possibly of other sequences surrounding the transcription initiation site (initiator) and can either be used isolated (minimal promoter) or linked to binding sites of transcription-activating elements, silencers or enhancers that may enhance or reduce transcription initiation rates, and which may function respective of developmental stage, or external or internal stimuli.

The **initiation site** is the position surrounding the first nucleotide which is part of the transcribed sequence, which is also defined as position +1. With respect to this site all other sequences of the gene and its controlling regions are numbered. Downstream sequences (i.e. further protein encoding sequences in the 3' direction) are denominated positive, while upstream sequences (mostly of the controlling regions in the 5' direction) are denominated negative.

A **minimal promoter** is a promoter consisting only of all basal elements needed for transcription initiation, such as a TATA-box and/or initiator.

An **enhancer** is a DNA-element which, when present in the neighbourhood of a promoter is able to increase the transcription initiation rate.

A promoter is **constitutive** when it is able to express the gene that it controls in all or nearly all of the plant tissues during all or nearly all developmental stages of the plant.

5 **Specific expression** is the expression of gene products which is limited to one or a few plant tissues (spatial limitation) and/or to one or a few plant developmental stages (temporal limitation). It is acknowledged that hardly a true specificity exists: promoters seem to be preferably switch on in some tissues, while in other tissues there
10 can be no or only little activity. This phenomenon is known as **leaky expression**. However, with specific expression in this invention is meant preferable expression in one or a few plant tissues.

15 The **expression pattern** of a promoter (with or without enhancer) is the pattern of expression levels which shows where in the plant and in what developmental stage transcription is initiated by said promoter.

20 Expression patterns of a set of promoters are said to be complementary when the expression pattern of one promoter shows little overlap with the expression pattern of the other promoter.

25 The level of expression of a promoter can be determined by measuring the 'steady state' concentration of a standard transcribed reporter mRNA. This measurement is indirect since the concentration of the reporter mRNA is dependent not only on its synthesis rate, but also on the rate with which the mRNA is degraded. Therefore the steady state level is the product of synthesis rates and degradation rates. The rate of degradation can however be considered to proceed at a fixed
30 rate when the transcribed sequences are identical, and thus this value can serve as a measure of synthesis rates. When promoters are compared in this way techniques available to those skilled in the art are hybridisation S1-RNase analysis, Northern blots and competitive RT-PCR. This list of techniques in no way represents all available
35 techniques, but rather describes commonly used procedures used to analyse transcription activity and expression levels of mRNA.

One of the technical difficulties encountered in such an analysis is that the qualitatively best results can only be obtained by fusing transcriptional activating parts to the reporter RNA molecule, in such

a way that only reporter sequences are transcribed. This requires the exact determination of the RNA synthesis start, and joining at that point the sequences of the reporter mRNA.

5 This is important for a number of reasons. First, the analysis of transcription start points in practically all promoters has revealed that there is usually no single base at which transcription starts, but rather a more or less clustered set of initiation sites, each of which accounts for some start points of the mRNA. Since this
10 distribution varies from promoter to promoter the sequences of the reporter mRNA in each of the populations would differ from each other. Since each mRNA species is more or less prone to degradation, no single degradation rate can be expected for different reporter mRNAs. Secondly, it has been shown for various eukaryotic promoter sequences
15 that the sequence surrounding the initiation site ('initiator') plays an important role in determining the level of RNA expression directed by that specific promoter. This includes also part of the transcribed sequences. The direct fusion of promoter to reporter sequences would therefore lead to much suboptimal levels of transcription.

20 Leaving in these transcribed sequences does allow determining the transcription rates, but potentially alters the stability of the reporter mRNA and influences translation initiation rates of an eventual open reading frame.

25 The role of this analysis, however, is the determination of the relative level of constitutive expression of a heterologous protein, as is the most frequent used application in biotechnology. Therefore the most important parameter is the ability of the tested sequences to
30 drive high level expression of a heterologous reporter protein.

This would involve coupling the coding sequences of a reporter protein to the transcription activating part, promoter and 5' untranslated sequence of the gene which is tested for its properties. In this way a
35 complex set of effects (combining transcription rates, mRNA stability (and thus degradation rates of the mRNA) and translational initiation rates) is reduced to one value that is a very useful value for determining usefulness of the tested gene elements in biotechnological applications.

There is no current word or phrase to describe this value. In the course of this application next to the term '**expression value**' the terms 'expression level' and 'transcriptional activity' are used. We realize that this may cause some confusion. In all cases we do

indicate with these and related terms the value just mentioned. A commonly used procedure to analyse expression patterns and levels is then through determination of the 'steady state' level of protein accumulation in a cell. Commonly used candidates for the reporter gene, known to those skilled in the art are β -glucuronidase (GUS), Chloramphenicol Acetyl Transferase (CAT) and proteins with fluorescent properties, such as Green Fluorescent Protein (GFP) from *Aequora victoria*. In principle, however, many more proteins are suitable for this purpose, provided the protein does not interfere with essential plant functions. For quantification and determination of localization a number of tools are suited. Detection systems can readily be created or are available which are based on e.g. immunochemical, enzymatic, fluorescent detection and quantification. Protein levels can be determined in plant tissue extracts or in intact tissue using in situ analysis of protein expression.

Generally, individual transformed lines with one chimeric promoter-reporter construct will vary in their levels of expression of the reporter gene. Also frequently observed is the phenomenon that such transformants do not express any detectable product (RNA or protein). The variability in expression is commonly ascribed to 'position effects' although the molecular mechanisms underlying this inactivity are usually not clear.

The term **average expression** is used here as the average level of expression found in all lines that do express detectable amounts of reporter gene, so leaving out of the analysis plants that do not express any detectable reporter mRNA or -protein.

Root expression level indicates the expression level found in protein extracts of complete plant roots. Likewise, 'leaf' and 'stem expression levels' are determined using whole extracts from leaves and stems. It is acknowledged however, that within each of the plant parts just described, cells with variable functions may exist, in which promoter activity may vary.

For the promoters described in this application the expression levels in large plant parts, containing cells with various functions, are measured. However, more detailed analyses may contribute to

construction of a promoter that is even 'more constitutive' taking into account that more celltypes within a plant part are taken into account.

As a standard for judging expression levels the 35S promoter of the Cauliflower Mosaic Virus is a convenient and widespread used standard. The average expression level of this promoter may be classified as medium high.

The invention shows that it is possible to combine elements from one promoter which are responsible for a specific expression with elements from another promoter which are responsible for a complementary expression pattern to form a promoter which - as a result - shows expression in the tissues and developmental stages which form part of the expression pattern of both promoters. If the complementation results in activity in (nearly) all the cells of the plant, such complementation will yield a constitutive promoter. It seems to be necessary, however, that both promoters have a low expression value in the tissues and developmental stages which are specific for the other promoter. It has been established that, for being suitable, the transcriptional activity in the plant parts where expression is low should be preferably $\geq 1\%$ of the level of transcription which is reached in the plant parts where transcriptional activity is high.

This limits the availability of promoters and promoter elements from which to build a new constitutive promoter. A suitable promoter-pairs which fulfills the above mentioned criteria is:

- the ferredoxin promoter in combination with the rolD promoter
- the S-adenosyl methionine promoter in combination with the plastocyanin promoter

Other promoter-pairs which are complementary and which show at least some expression in the tissues and developmental stages which are specific for the other promoter can also be applied.

Delineation of promoter and/or enhancer parts needed.

Whereas transcription-regulating elements, especially in eukaryotes, may be present at large distances from the promoter/transcription initiation site, and located both downstream or upstream of the initiation site, many plant genes have most of their regulatory elements in the area directly upstream of the promoter. In order to

identify the main transcription-activating elements of promoters it is common procedure to link parts of the non-transcribed areas that are found upstream (and downstream) of the promoter to a reporter gene, to analyse the ability of each of the truncated DNA elements to direct expression of that reporter. For delineation of more promoter-proximal sequences involved in transcription regulation, fragments of the enhancer sequences are most commonly coupled to a promoter, which may be derived from the gene of which transcription regulation is studied. Alternatively, a heterologous promoter can be used such as the sequences of the 35S promoter from -46 to +4, relative to the transcription start, which is functionally coupled to a reporter gene as described above.

In this way it is possible to delineate the transcription activating elements of most genes, a process that is well-known to those skilled in the art.

A large number of transcription regulatory elements of genes have been analysed in such a manner, and data relevant for this analysis are directly available to those skilled in the art through scientific publications.

Transcription activating elements that on average can direct expression to approximately the average level of the 35S promoter (at least 50% of this level) in at least some of the plant parts, and that are also capable of directing at least 0.5% (of the 35S level) transcription in other plant parts are then selected for further use.

The minimal promoter element is typically derived from one of the promoters of the promoter-pair, although not necessarily. It can be envisaged that such a minimal element is derived from a third promoter or is even made synthetically.

Based on the results of the analysis described above, transcription activating parts with complementary activities are selected. That is, for example, a promoter with expression throughout the plant, transcription activating DNA fragments that direct high level root expression and with lower leaf and stem expression levels, are combined with elements that direct expression mainly in the leaf and stem, but lower in the root. Other combinations of complementary transcription activating parts are obvious.

Preferentially, the level of expression in the parts where expression is lowest does not fall below 1% of the level obtained in the highest

part. More preferred is the situation where the relation between lowest expression and highest expression between plant parts is larger than 5%.

This coupling can most easily be done by known genetic engineering techniques. The gene which has to be expressed by the new constitutive promoter can be cloned behind the promoter. It is advisable to build in a unique NcoI-cloning site at the linkage of the 5' untranslated sequence attached to the promoter to allow precise junction of the open reading frame (ORF) and the 3' end of the promoter in which the gene of interest can be inserted.

The ferredoxin-rolD pair.

One of the preferred combinations of the present invention is a constitutive plant promoter comprising elements of both the ferredoxin promoter and the rolD-promoter. Preferably the ferredoxin promoter is obtained from *Arabidopsis thaliana* where it drives the ferredoxin A gene, a gene which is involved in the photosynthesis. The expression of this gene and the responsiveness of its promoter to light has been reported (Vorst, O. et al., Plant Mol. Biol. 14, 491-499, 1990; Vorst, O. et al., The Plant J. 3(6), 793-803, 1993; Dickey, L.F. et al., The Plant Cell 6, 1171-1176, 1994). Since the ferredoxin gene is involved in photosynthesis the promoter is most active in green tissue. mRNA levels were shown to be high in chloroplast-containing organs such as stem, leaves and bracts, but also in young growing tissues, such as whole flowers and seedlings. Interestingly, there is a smaller, but significant expression in soilborne areas of the plant. The promoter sequence contains both a G-box and an I-box containing region. Also a potential Z-folding DNA sequence is found at position -182.

The rolD promoter is reported to have strong expression in the roots and is obtainable from *Agrobacterium rhizogenes*. Although the source organism is a bacterium, the promoter is very suitable for expression in plants because the bacterium is a phytopathogen which causes hairy-root disease in plants. For that purpose it transfers DNA to the plant amongst which the rolD gene is responsible for root elongation. To be expressable in plants this gene needed a strong promoter functional in plants, the rolD promoter. GUS-studies have shown that expression under control of the rolD-promoter yields mainly root-specificity (Leach, F. and Aoyagi, K., Plant Sci. 79, 69-76, 1991). Also, some expression in leaves was observed.

A combination of the ferredoxin and the rolD promoter can be obtained in two ways, depending on from which promoter the minimal promoter element and 5' untranslated sequences will be taken. In our examples we have used the minimal promoter element from the ferredoxin promoter, but deriving it from the rolD promoter is equally well possible.

The S-adenosyl-methionine synthetase and plastocyanin pair.

Another favorable promoter can be obtained from a combination of the S-adenosyl-methionine synthetase (SAM) promoter and specific parts of the plastocyanin promoter. Preferably, both promoters are obtained from *Arabidopsis thaliana*.

The SAM promoter regulates the expression of S-adenosyl-methionine synthetase, which is an enzyme active in the synthesis of polyamines and ethylene. Promoter studies showed a strong expression in vascular tissues, in callus, sclerenchyma and some activity in root cortex (Peleman, J. et al., The Plant Cell 1, 81-93, 1989) which was reasoned to be due to the involvement of the enzyme in lignification.

The plastocyanin promoter, like the ferredoxin promoter, is also a promoter which is active in the photosynthetic pathway. mRNA levels are high in green, chloroplast-containing structures, such as leaves, cauline leaves, stem and whole seedling. Also in flowers the promoter is very active. Little expression is detectable in silique, seed and root (Vorst, O. et al., The Plant J. 4(6), 933-945, 1993).

By combining these specificities it is possible to create a chimeric promoter that drives good expression both in the photosynthetic areas of the leaf and stem, as well as in the area's not involved in photosynthesis, such as the cells forming and surrounding the vascular system in leaves and stems.

Other pairs of promoters.

The above given examples of promoter-pairs show in both cases the presence of a promoter which is active during photosynthesis. It is envisaged that other promoters which are regulating expression of a gene needed for photosynthetic activity may be suitable for a combination with either the rolD or other root-preferential promoters.

In the construction of a promoter that drives expression throughout the plant: if one of the components is a promoter which is more or less specific for green parts, this automatically means that

the other promoter of the pair should be predominantly (but not exclusively) expressed in the roots and other non-photosynthesizing organs.

5 In the construction of a promoter that drives expression in all parts of leaves and stems, the combination may be made by using a promoter which is more or less specific for green parts and a promoter which drives expression primarily in the vascular system.

10 However, the invention is not limited to the combination of a root-preferential and a green part-preferential promoter, and a combination of green-part-preferential and vascular system-preferential promoters. All promoter combinations provided that the expression patterns of the individual promoters are complementary can be used.

15 It is also possible that the elements from which e.g. a new constitutive promoter is composed are derived from a set with more than two promoters. The above discussed complementarity should then also exist.

EXPERIMENTAL PART

Example 1

Cloning of the chimeric Fd-rolD promoter:

5 A 512 bp *Arabidopsis thaliana* ferredoxin promoter fragment (O. Vorst et al., 1990, PMB 14, 491-499.) ranging from position -512 to +4 (relative to the ATG startcodon of the ferredoxin Open Reading Frame) was isolated by digestion with HincII and NcoI. This fragment contains most of the transcriptional regulatory sequences of the ferredoxin promoter, the promoter sequences and leader of the ferredoxin transcript. An XbaI site was introduced, for cloning reasons, at positions -5 to -10 relative to the ATG (O. Vorst et al., 1990, PMB 14, 491-499.). This changes the original sequence of the clone at this point from ACAAAA to TCTAGA (SEQ ID NO: 1).

10 Part of the *Agrobacterium rhizogenes* rolD upstream sequences (SEQ ID NO: 2) (Leach et al., 1991 Plant Sci. 79, 69-76) were fused to the ferredoxin promoter sequences described above. A HindIII-RsaI fragment, comprising nucleotides -385 to -86 relative to the initiation codon was cloned next to the ferredoxin fragment, joining the RsaI sites of the latter with the HincII site of the former. This chimaeric element, containing the promoter and some of the activating sequences of the ferredoxin gene, and upstream activating sequences of the rolD gene was used in subsequent studies as to its transcription-stimulating properties (SEQ ID NO: 3).

15

20

25

Example 2

GUS-fusions

The Fd-rolD chimaeric promoter/activator was coupled to the GUS gene, engineered to contain an intron gene (Jefferson et al., (1987) EMBO J 6: 3901-3907). The NcoI restriction site on the ATG start codon was used to join the promoter to the Open Reading Frame (ORF) of the GUS gene, coupled to a 265 bp fragment containing the Proteinase Inhibitor II 3' untranslated and transcriptional termination sequences (Thornburg et al., 1987, Proc. Natl. Acad. Sci. USA 84, 744-748; An et al., Plant Cell 1, 115-122).

30

35

The whole expression cassette, containing the promoter, GUS gene and 3' PI-II sequences was cloned out using BamHI and EcoRI and introduced into the binary vector pMOG800 (deposited at the Centraal Bureau voor

Schimmelcultures, Baarn, The Netherlands, under CBS 414.93, on august 12, 1993) digested with the same enzymes. The subsequently made construct (pMOG1059) was used in transformation experiments with various plants. As a control a 35S CaMV promoter-GUS construct was used. This is construct pMOG410. A schematic representation of both constructs is found in figure 1.

Example 3

Expression levels and patterns of promoter activity during early stages of plant transformation

First, *Arabidopsis thaliana* transformants were made with both constructs and GUS expression was followed in time during the transformation procedure.

GUS expression levels were determined visually, on a scale of 0 to 5, where 0 is no detectable expression and 5 is the highest level of GUS we have observed in leaves of a transgenic plant, of a rare tobacco 35S-GUS-transgenic (line 96306). Samples from leaves of this plant were included in all experiments for internal reference.

In table 1 the relative GUS expression in *Arabidopsis thaliana* explants is indicated, at several times after *Agrobacterium tumefaciens* cocultivation (DAC; days after cocultivation)

Table 1. relative GUS activity of *Arabidopsis* root explants.

Construct :	pMOG1059	pMOG410
Time of assay		
DAC 0	2	3
DAC 2	3	3
DAC 5	3	3
DAC 7	4	3
DAC 9	4	3
DAC 12	4	3

As can be seen from this comparison, GUS expression driven by the chimaeric promoter starts slightly later after cocultivation but from day 7 on, exceeds the level of expression obtained with the reference 35S promoter.

Very similar data were obtained when *Brassica napus* explants were scored for GUS expression. At day 5 after co-cultivation the 35S promoter is slightly higher, but the situation is reversed on day 20 after co-cultivation. Also for tomato similar data were obtained. Here

even at the earliest stage of analysis expression of pMOG1059-transgenics exceeded that of pMOG410 transgenics.

Example 4

Expression levels and patterns in in vitro grown plants

When plants are grown up further, differences between these promoters become ever clearer. Leaf samples of fully regenerated plants were analysed for GUS expression. Averages were obtained from 11-48 plants, dependent on the construct.

For *Arabidopsis thaliana* that was grown in vitro only, no large difference was observed between GUS expression in pMOG1059 and pMOG410-transgenics.

Table 2. Average relative GUS activity of leaf samples of all tested crops.

Construct :	pMOG1059	pMOG410
Crop :		
Potato	4.0	2.1
Brassica napus	3.7	2.8
Arabidopsis	4.0	4.1
Tomato	2.2	2.1

What is also clear from the data presented in figure 2 that a significant number of 35S-GUS transgenic lines (app. 50% was found repeatedly in our experiments) do not express GUS to a level that it is visible. So not only maximum and average expression are higher in the Fd-rolD-GUS transgenics, also the frequency with which transgenic plants do express GUS is strongly enhanced. In about 50 transgenic potato plants carrying the Fd-rolD-GUS construct, we have found no weak expressor, suggesting a reliable high expression in at least 98% of the lines made.

Example 5

Comparison of promoter performance in various crops

Constructs pMOG410 (35S-GUS) and pMOG1059 (Fd-rolD-GUS) were also introduced into oilseed rape and tomato for a further comparison of promoter performance. Also the data for potato are included here.

As shown in Figure 3A, in tomato the overall level of expression of the Fd-rolD promoter is higher both at the latest stage of in vitro growth as well as in leaves of 4 and 7 week old plants. Also in stems of 7 weeks this holds true, however, for roots, an average weaker expression is observed with the Fd-rolD promoter than for the 35S promoter.

Also in oilseed rape and potato, similar results are obtained, with the notable exception that in potato roots the level of expression by the Fd-rolD promoter exceeds that of the 35S promoter. As shown in figures 3B and 3C both the average expression of the Fd-rolD promoter is higher and also the variation in expression is significantly lower. In conclusion we can say we have created a promoter that withstands the comparison with the 35S promoter easily in three major crops.

Example 6

Expression of nptII transgene.

In order to also check usability of the Fd-rolD promoter for other purposes, the promoter was linked to the nptII gene, of which expression of the corresponding gene product confers resistance in plants to the antibiotic kanamycin. This element was placed between the left and right borders of the T-DNA allowing *Agrobacterium tumefaciens*-mediated transfer to plants. As a control, similar constructs in which the expression of the nptII gene was under control of the nos promoter were used.

The resistance to kanamycin in transgenic potato plants is manifested by the development of transgenic calli and shoots during a standard transformation procedure, in which kanamycin is used in the culture medium.

On average, for the constructs with the nos-nptII selection cassette, the transformation frequency for potato is 45%, for constructs with the Fd-rolD-nptII selection cassette the frequency is on average 61%. While we do not know at this moment how relevant the increase in transformation frequency is for this construct, it indicates that the Fd-rolD promoter is at least as suitable for driving a heterologous gene such as nptII, as commonly used constitutive promoters such as nos.

Example 7**Comparing visual scoring to quantitative values**

From the analysis of GUS expression based on 1) histochemical analysis and scoring to an internal control and 2) to a quantitative analysis of GUS enzymatic activity, we have learned that both give a reproducible quantitative figure. A thorough analysis of both scores for tomato and oilseed rape leaves and roots, leads to the conclusion that scale 3, which compares best to that of 35S, equals about 2000 pmol MU/minute.mg made in the quantitative analysis. In scale 1 and 2, averages are 1000 and 1500, respectively, which set the value of 50% of 35S. About 1% expression of that level equals 100 pmol MU/minute.mg, which is frequently under the detection level for histochemical detection, although sometimes detectable as very light blue staining due to GUS expression. Therefore one can use histochemical staining as a marker for promoter efficacy, by measuring the level of blue staining, and use these data to select promoter elements of use.

Example 8**Construction of the SAM1 promoter and fusion to GUS.**

For the construction of the SAM1 promoter genomic DNA (SEQ ID NO: 4) was isolated from *Arabidopsis thaliana* Landsberg erecta leaves using a CTAB extraction procedure. Primers were designed based on the published sequence of the SAM1 gene from *Arabidopsis thaliana* K85 (Peleman et al., (1989) Gene 84, 359-369). In a PCR (30 cycles of 45 seconds 95 °C, 45 seconds 50°C and 1' 72°C; same program was used in all other PCR's described in this part) the promoter element was amplified using primers FR-Psam-143 5' AGA TTT GTA TTG CAG CGA TTT CAT TTT AG 3' (SEQ ID NO: 5) and FR-Psam-216 5' ATC TGG TCA CAG AGC TTG TC 3' (SEQ ID NO: 6) yielding a fragment of about 550 bp. The DNA fragment was isolated from an agarose gel and cloned into the pGEM-T vector (Promega Corp., Madison WI, USA). This clone was used as a template to introduce a Nco I site at the translation start by PCR using primers FR-Psam-144 5' GTC TCC ATG GTG CTA CAA AGA ATA G 3' (SEQ ID NO: 7) and FR-Psam-143. The resulting 500 bp fragment was cloned in the pGEM-T vector. The EcoR I and Hind III sites located in the promoter region were removed by PCR in two steps using this clone

as a template. In this PCR a *Bam*H I site was introduced upstream of the SAM1 promoter and a *Hind* III site was introduced at the 3' site of the promoter. In the first PCR step three promoter fragments were generated. The first fragment (1) will contain the 5' *Bam*H I site and the mutated *Eco*R I site using primers FR-Psam-248 5' CGG GAT CCT GCA GCG ATT TCA TTT TAG 3' (SEQ ID NO: 8) and FR-Psam-249 5' ACA TGA ACG AAT GCA AAA TCT C 3' (SEQ ID NO: 9). The middle fragment (2) is obtained with primers FR-Psam-250 5' AGA TTT TGC ATT CGT TCA TGT G 3' (SEQ ID NO: 10) and FR-Psam-251 5' TGT AAG CAT TTC TTA GAT TCT C 3' (SEQ ID NO: 11). This fragment has a partial overlap with fragment 1 and 3 and has mutated *Eco*R I and *Hind* III sites. The third PCR fragment (3) will contain the mutated internal *Hind* III site and introduces a *Hind* III site at the 3' end of the promoter encompassing the *Nco* I site at the translation start and is generated using primers FR-Psam-252 5' AAG AAA TGC TTA CAG GAT ATG G 3' (SEQ ID NO: 12) and FR-Psam-253 5' GAC AAG CTT GAT CCC ATG GTG CTA CAA AGA ATA G 3' (SEQ ID NO: 13). In a second PCR the 3 fragments 1, 2 and 3 were mixed together in one tube and amplified with primers FR-Psam-248 and FR-Psam-253. Due to the overlap between fragments 1 and 2, and 2 and 3, this PCR yields the complete mutated promoter. After digestion with *Bam*H I and *Hind* III the resulting SAM1 promoter was cloned in a pBSK+ vector. The SAM1 promoter was then cloned into a vector containing a GUSintron-TPI-II reporter cassette by exchanging the upstream region using the *Bam*H I and *Nco* I restriction sites. This was done by digestion of the SAM1 clone with *Bam*H I and *Nco* I and isolation of the promoter fragment from a agarose gel. The GUS vector was digested with the same enzymes and the vector was then isolated from a agarose gel thus discarding the original upstream sequences promoter.

The SAM1 promoter-GUSintron-TPI-II reporter cassette was then cut out of the vector by *Bam*H I and *Eco*R I digestion after which the reporter cassette was isolated from a agarose gel and cloned into the binary vector pMOG800 digested with *Bam*H I and *Eco*R I. The resulting binary vector pMOG1402 was introduced in *Agrobacterium tumefaciens* strain EHA105 for transformation to potato.

Example 9**Construction of the Pc-SAM1 chimaeric promoter and fusion to the GUS gene**

5 The plastocyanin enhancer (Pc) from *Arabidopsis thaliana* Col-0 was obtained by PCR. Therefore primer FR-Pc-146 (5' agt ggt acc atc ata ata ctc atc ctc ctt ca3') (SEQ ID NO: 14) and primer FR-Pc-247 (5' cga agc ttt aca aat cta att tca tca cta aat cgg a3') (SEQ ID NO: 15) were developed introducing a *Kpn* I restriction site upstream of the
 10 enhancer and a *Hind* III restriction site downstream of the plastocyanin enhancer. The PCR was performed using Cloned *pfu* DNA polymerase (Stratagene) for 30 cycles 1' 95°C, 1' 50°C, 4' 72°C and 1 cycle 1' 95°C, 1' 50°C, 10' 72°C. The resulting PCR fragment was ligated into a high copy cloning vector using *Kpn* I and *Hind* III
 15 resulting in construct pPM15.1. This clone was used as a template for a PCR (30 cycles of 1' 95 °C, 1' 50°C and 2' 72°C) using primers FR-Pc-145 5' GCT GCA ATA CAA ATC TAA TTT CAT CAC TAA ATC GG 3' (SEQ ID NO: 16) and FR-Pc-146 5' AGT GGT ACC ATC ATA ATA CTC ATC CTC CTT C 3' (SEQ ID NO: 14). The PCR generates a
 20 fragment of about 850 bp encompassing the Pc enhancer (SEQ ID NO: 17) containing a upstream *Kpn* I site and overlap with the 5' side of the SAM1 promoter (see Example 8). The PCR fragment was then mixed with a PCR fragment of the SAM1 promoter generated with primers FR-Psam-143 and FR-Psam-144 using the pBKS+ clone containing the adjusted SAM1
 25 promoter described in example 8. In a PCR on this mixture the PcSAM chimeric promoter was generated using primers FR-Psam-144 and FR-Pc-146. The resulting promoter fragment of about 1.3 kb (SEQ ID NO: 18) was isolated from a agarose gel after digestion with *Kpn* I and *Nco* I and then cloned into a high copy cloning vector (pUC28) digested with
 30 the same enzymes. The promoter fragment was then cut out of this vector by digestion with *Bam*HI and *Nco*I and cloned in front of the GUSintron gene as described above in Example 8. The complete Pc-SAM-GUS-TPI-II reporter cassette was then cloned into pMOG800 as described for the SAM1-GUS-TPI-II reporter cassette in Example 8. The resulting
 35 binary vector pMOG1400 was introduced in *Agrobacterium tumefaciens* strain EHA105 for transformation to potato.

Example 10**Construction of the Pc enhancer-35S promoter and fusion to the GUS gene**

5 The plastocyanin enhancer (Pc) from *Arabidopsis thaliana* Col-0 was obtained by PCR (see above).

This clone was used as a template for a PCR (30 cycles of 1' 95 °C, 1' 50°C and 2' 72°C; all other PCR reactions described in this part were carried out with the same program) using primers FR-Pc-291 5' GTC TTG TAC AAA TCT AAT TTC ATC ACT AAA TCG G 3' (SEQ ID NO: 19) and FR-Pc-146 10 5' AGT GGT ACC ATC ATA ATA CTC ATC CTC CTT C 3' (SEQ ID NO: 14). The PCR generates a fragment of about 850 bp encompassing the Pc enhancer containing a upstream KpnI site and overlap with the 5' side of the minimal 35S promoter. The minimal 35S promoter was obtained in a PCR 15 using pMOG971 as a template (containing the 35S promoter and omega 5' UTR) and primers FR-35S-292 5' TTA GAT TTG TAC AAG ACC CTT CCT CTA TAT AAG G 3' (SEQ ID NO: 20) and ls19 (SEQ ID NO: 21). The resulting fragment has overlap with the Pc enhancer and contains a internal NcoI site at the translation start. The two PCR fragments were then mixed 20 and a PCR reaction was carried out using primers FR-Pc-146 and ls19. The resulting fragment was then digested with KpnI and NcoI, isolated from a agarose gel and cloned in pUC28 digested with the same enzymes. The resulting clone was, subsequently, digested with BamHI and NcoI, and the promoter fragment (SEQ ID NO: 22) was isolated from a agarose 25 gel and cloned upstream of the GUS gene as described in Example 7. The complete reporter cassette was then introduced in the binary vector pMOG800 as described in Example 7. The resulting binary vector pMOG1401 was introduced in *Agrobacterium tumefaciens* strain EHA105 for transformation to potato.

Example 11**Expression levels and patterns in in vitro grown plants**

Transformed plants were grown up, and leaf samples of fully 35 regenerated plants were analyzed for GUS expression. In figure 4 the analysis of expression in leaf mesophyll, leaf vascular system, stems and roots is indicated. A very low level of GUS staining was observed in the mesophyll part of leaves of SAM1-transgenic plants although the scoring indicates a GUS expression level of 0.

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

Mogen International N.V.
Einsteinweg 97
2333 CB LEIDEN
Nederland

name and address of depositor

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the
DEPOSITOR:

E. coli DH5 alpha strain / the plasmid
pMOG800

Accession number given by the
INTERNATIONAL DEPOSITARY AUTHORITY:

CBS 414.93

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:



a scientific description



a proposed taxonomic designation

(mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary accepts the microorganism identified under I above, which was
received by it on Thursday, 12 August 1993 (date of the original deposit)¹

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority on not applicable (date of the original deposit) and a
request to convert the original deposit to a deposit under the Budapest Treaty was received by
it on not applicable (date of receipt of request for conversion)

V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Centraalbureau voor Schimmelcultures

Address: Oosterstraat 1
P.O. Box 273
3740 AG BAARN
The Netherlands

Signature(s) of person(s) having the power to
represent the International Depositary
Authority or of authorized official(s):



drs F.M. van Asma

Date: Friday, 13 August 1993

¹ Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired.

AMENDED CLAIMS

1. Chimaeric plant promoter, characterized in that it comprises a minimal promoter and transcription-activating elements from a set of promoters, which elements have a complementary pattern and level of transcription in a plant.
2. Chimaeric plant promoter according to claim 1, characterized in that each of the transcription-activating elements do not exhibit an absolute tissue-specificity, but mediate transcriptional activation in most plant parts at a level of $\geq 1\%$ of the level reached in the part of the plant in which transcription is most active.
3. Chimaeric plant promoter according to claim 1 or 2, characterized in that one promoter of the set of promoters is specifically active in green parts of the plant, while the other promoter is specifically active in underground parts of the plant.
4. Chimaeric plant promoter according to claim 3, characterized in that it is a combination of the ferredoxine and the Rold promoter.
5. Chimaeric plant promoter of claim 4, characterized in that the minimal promoter element is derived from the ferredoxin promoter.
6. Chimaeric plant promoter according to claim 4 or 5, characterized in that the ferredoxin promoter is derived from *Arabidopsis thaliana*.
7. Chimaeric plant promoter according to claim 6, characterized in that it comprises the sequences of SEQ ID NO: 1 and SEQ ID NO: 2.
8. Chimaeric promoter according to claim 7, characterized in that it comprises the sequence of SEQ ID NO: 3.

AMENDED SHEET

9. Chimaeric plant promoter according to claim 3,
characterized in that it is a combination of the plastocyanin
and the S-adenosyl-methionine-1 promoter.
10. Chimaeric plant promoter according to claim 9,
characterized in that the minimal promoter element is derived
from the S-adenosyl-methionine-1 promoter.
11. Chimaeric plant promoter according to claim 9 or 10,
characterized in that the plastocyanin promoter is derived from
Arabidopsis thaliana.
12. Chimaeric plant promoter according to claim 9, 10 or 11,
characterized in that the S-adenosyl-methionine-1 promoter is
derived from *Arabidopsis thaliana*.
13. Chimaeric plant promoter according to claim 12,
characterized in that it comprises the sequences of SEQ ID NO:4
and SEQ ID NO:17.
14. Chimaeric plant promoter according to claim 13,
characterized in that it comprises the sequence of SEQ ID NO:
21.
15. Chimaeric gene construct for the expression of genes in
plants comprising the promoter of any of claims 1-14.






Fig. 1


Fig. 2




Tomato




pMOG1059




Brassica



pMOG1059



Potato



pMOG1059





Fig. 3



Sheet1 Chart 3

Fig. 4

Docket No. MOG57701/UST

- ☐ Declaration Submitted with Initial Filing
☒ Declaration Submitted after Initial Filing (surcharge (37 CFR 1.16(e)) Required)

DECLARATION FOR UTILITY OR DESIGN PATENT APPLICATION (37 CFR 1.63)

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

CONSTITUTIVE PLANT PROMOTERS

the specification of which

☐ is attached hereto

OR

☒ was filed on December 10, 1998 as United States Application Number **OR** PCT International Application Number PCT/EP98/08162 and was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT International application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
97203912.7	EP	December 12, 1997	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

I hereby claim the benefit under 35 U.S.C. §119(e) of any United States **provisional** application(s) listed below.

Provisional Application Number(s)	Filing Date (MM/DD/YYYY)

I hereby claim the benefit under 35 U.S.C. §120 of any United States application(s), **or** 365(c) international application designating the United States of America listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose information which is material to patentability as defined in 35 CFR 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application or PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)

4- I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:
Dianne Burkhard, Reg. No. 41,650; Liza D. Hohenschutz, Reg. No. 33,712; Thomas R. Savitsky, Reg. No. 31,661; William E. Dickheiser, Reg. No. 30,169.

Address all telephone calls to Liza D. Hohenschutz at telephone number **(302)886-1699**.

Address all correspondence to: ZENECA Ag Products Inc.
Intellectual Property Department
1800 Concord Pike
P.O. Box 15458
Wilmington, DE 19850-5458

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

5-00

Name of Sole or First Inventor:							
Given Name (first and middle [if any])				Family Name or Surname			
Maarten Hendrik				STUIVER			
Inventor's Signature				Date		26-06-00	
Residence:	Oegstgeest	STATE		COUNTRY	The Netherlands	Citizenship	The Netherlands
CITY				NLX			
Post Office Address:		Groenhoevelaan 71					
City	Oegstgeest	State		ZIP	2343 BR	Country	The Netherlands

2-00

Name Second Inventor:							
Given Name (first and middle [if any])				Family Name or Surname			
Floor Hendrik				SIJBOLTS			
Inventor's Signature				Date		18-06-2000	
Residence:	Almere	STATE		COUNTRY	The Netherlands	Citizenship	The Netherlands
CITY				NLX			
Post Office Address:		Pastelstraat 144					
City	Almere	State		ZIP	1339 JC	Country	The Netherlands

Name of Third Inventor:							
Given Name (first and middle [if any])				Family Name or Surname			
Inventor's Signature				Date			
Residence:		STATE		COUNTRY		Citizenship	
CITY							
Post Office Address:							
City		State		ZIP		Country	

SEQUENCE LISTING

5 (1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: Zeneca MOGEN
 (B) STREET: Einsteinweg 97
 (C) CITY: Leiden
 (E) COUNTRY: The Netherlands
 (F) POSTAL CODE (ZIP): 2333 CB
 (G) TELEPHONE: (31) 71-5258282
 (H) TELEFAX: (31) 71-5221471

(ii) TITLE OF INVENTION: New constitutive plant promoters

(iii) NUMBER OF SEQUENCES: 22

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
 (B) COMPUTER: IBM PC compatible
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

(vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: EP 97203912.7
 (B) FILING DATE: 12-DEC-1997

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 520 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GACTGAAGTG TGAAGGTGGA GATTATGTAT TCACTTGTTG ATTTGGTATA CATTCTATGT	60
AAGGTTCAAT TATTTACGTT ATATAATTAT AATGGAGTAA TTTACAGTAA TTGGGTAA	120
ATGGTTTGAT TCGGTCAGGT TGATACGGTT TGGAAGTTAA ACCCGGCCTA GATATGATGT	180
TACAACCACT CCACATCTTT TATGATTTTA GTGGAACAAA CGAAGAGTTA TTTAGACGAT	240
ACAAACAAGG TCCGAATAAG TGTGAGCTGT CCCAAGTAAG ACCACGTAAT ACTCACCTCA	300
ACAAGATAGT GTTCTTAAAG TGTGTCAAAC ACAATCACAC ACACACAAAT CATAAAACAC	360
AAAGACGATA ATCCATCGAT CCACAGAATA GACGCCACGT GGTAGATAGG ATTCTCACTA	420
AAAAGTTCTC ACCTTTTAAAT CTTTCTCCAC GCCATTTCCA CAAGCCATAA TCCTCAAAAA	480
TCTCAACTTT ATCTCCCAA ACACAAATCT AGAAACCATG	520

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 300 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CCCACTACAA TGAATTTGTT CGTGAACATAT TAGTTGCGGG CCTTGGCATC CGACTACCTC 60
 TGGCGCAATA TTATATTCCC TGGGCCACCC GTGAACCCAA TTTCGCCTAT TTATTCATTA 120
 CCCCCATTAA CATTGAAGTA GTCATGATGG GCCTGCAGCA CGTTGGTGAG GCTGGCACAA 180
 CTCATCCATA TACTTTCTGA CCGGATCGGC ACATTATTGT AGAAAACGCG GACCCACAGC 240
 GCACTTTCCA AAGCGGTGCC GCGTCAGAAT GCGCTGGCAG AAAAAAATTA ATCCAAAAGT 300

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 840 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GGATCCGAGC TTGCATGCCC CCACTACAAT GAATTTGTTT GTGAACATAT AGTTGCGGGC 60
 CTTGGCATCC GACTACCTCT GCGGCAATAT TATATTCCCT GGGCCCACCG TGAACCCAAT 120
 TTCGCCTATT TATTCATTAC CCCCATTAACT ATTGAAGTAG TCATGATGGG CCTGCAGCAC 180
 GTTGGTGAGG CTGGCACAAC TCATCCATAT ACTTTCTGAC CGGATCGGCA CATTATTGTA 240
 GAAAACGCGG ACCCAGACG CACTTTCCAA AGCGGTGCCG CGTCAGAAATG CGCTGGCAGA 300
 AAAAAATTAA TCCAAAAGTG ACTGAAGTGT GAAGGTGGAG ATTATGTATT CACTTGTGTA 360
 TTTGGTATAC ATTCTATGTA AGGTTCAATT ATTTACGTTA TATAATTATA ATGGAGTAAT 420
 TTACAGTAAT TGGGTTAAAA TGTTTGTATT CGGTCAGGTT GATACGGTTT GGAAGTTAAA 480
 CCGGCGCTAG ATATGATGTT ACAACCACTC CACATCTTTT ATGATTTTAG TGGAACAAAC 540

GAAGAGTTAT TTAGACGATA CAAACAAGGT CCGAATAAGT GTGAGCTGTC CCAAGTAAGA 600
 CCACGTAATA CTCACCTCAA CAAGATAGTG TTCTTAAAGT GTGTCAAACA CAATCACACA 660
 5 CACACAAATC ATAAACACA AAGACGATAA TCCATCGATC CACAGAATAG ACGCCACGTG 720
 GTAGATAGGA TTCTCACTAA AAGATTCTCA CCTTTTAATC TTTCTCCACG CCATTTCAC 780
 AAGCCATAAT CCTCAAAAAT CTCAACTTTA TCTCCCAAAA CACAAATCTA GAAACCATGG 840
 10

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 477 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM:
- Arabidopsis thaliana*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

GGATCCTGCA GCGATTTTCAT TTAGATTCT CAAAAATATT CTCAGATGTG TGGGATTGA 60
 GTAGAGTTTA TGTTCGTTG GCATGATTG AATAGTATGC AAGATTTTGG AGATTTTGCA 120
 35 TTCGTTTCATG TGTGTATGTG TGATTGTAGC TTGATATGAT TTAACCTGTT AGTTAAATGT 180
 GCATAGACAA TAAGTAACAT ACGAAGCGAG TCACTAAGCA TAAGAGTCAA CTTGTTTTGC 240
 40 TGAAAAGATA TCACTTATGA TTTTCGAATC ATTTTAGCTT TTTGTCACT TGAGCTTAAT 300
 GATTCTTCTG AAATTCGATT CTTTGTGTTGG TTTATGTCAC ATTCTTTAGA ATTGAGAATC 360
 TAAGAAATGC TTACAGGATA TGGTGAAACT ATTCTTTTAA GATAGCATGA TGCTTCTTTT 420
 45 ATGATTCTAC AGTGGCTAAG TCATTTTTTT TTTGTTCTAT TCTTTGTAGC ACCATGG 477

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

AGATTTGTAT TGCAGCGATT TCATTTTAG 29
 65

(2) INFORMATION FOR SEQ ID NO: 6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

ATCTGGTCAC AGAGCTTGTC

20

(2) INFORMATION FOR SEQ ID NO: 7:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 25 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

GTCTCCATGG TGCTACAAAG AATAG

25

(2) INFORMATION FOR SEQ ID NO: 8:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

CGGGATCCTG CAGCGATTTC ATTTTAG

27

(2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 9:
5 ACATGAACGA ATGCAAAATC TC 22

(2) INFORMATION FOR SEQ ID NO: 10:
10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
15 (ii) MOLECULE TYPE: cDNA
 (iii) HYPOTHETICAL: NO

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:
25 AGATTTTGCA TTCGTCATG TG 22

(2) INFORMATION FOR SEQ ID NO: 11:
30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
35 (ii) MOLECULE TYPE: cDNA
 (iii) HYPOTHETICAL: NO

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:
TGTAAGCATT TCTTAGATTC TC 22

(2) INFORMATION FOR SEQ ID NO: 12:
45 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
50 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: cDNA
 (iii) HYPOTHETICAL: NO
55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:
60 AAGAAATGCT TACAGGATAT GG 22

(2) INFORMATION FOR SEQ ID NO: 13:
65 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 34 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

GACAAGCTTG ATCCCATGGT GCTACAAAGA ATAG

34

15 (2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

25 (iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

30

AGTGGTACCA TCATAATACT CATCCTCCTT C

31

(2) INFORMATION FOR SEQ ID NO: 15:

35 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 37 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

CGAAGCTTTA CAAATCTAAT TTCATCACTA AATCGGA

37

50

(2) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 35 base pairs

55 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

60

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

65 GCTGCAATAC AAATCTAATT TCATCACTAA ATCGG

35

(2) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 876 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Arabidopsis thaliana*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

ATCATAATAC TCATCCTCCT TCTCAAGGTT CGTACGTATT ATCAATATCT AGTATATACT 60
 TGTCTTTGTT CTATGCTTTA TATCATCATT TTATGACAAA AAATGATTAA GGTCTTAGTT 120
 AATGATTATG TATATGTGAA ACTTATATTT AGGGGCACAA TTTAATTTTCG TATGATAATT 180
 GTCTAGTTAG CTTTATGTAC TTATCATAAA AACCTTAGTG TTTATCGCAA TACTTTTCAA 240
 ATATAGTGTA GAATCATAAT GGTCCCACTG TCATTATGTT TGATGCAAAT CTATTTGGAT 300
 TTTGTTGGAT AATAAACCGA TGACGTGGAC CAGACCAGTA GCTATAAGAT TTGGTTCACA 360
 TAGAAATTTT TTATAAGATA ATGTATCTAG GTTTGCTTAT GATTATACAT GTGATATTTA 420
 ATACATGGCA CAGGTTTCGT CAGTTTCACA GCCATAGGTA CAATAGAAGG CAAATTCGAT 480
 TGTGGTTATC TGGTAAAAGT TAAGTTGGGC TCAGAGATTC TTAACGGCGT TCTTTATCAT 540
 TCGGCCAGC CCGGCCCATC ATCATCTCCA ACCGCTGTTC TAAACAATGC CGTTGTACCT 600
 TATGTTGAAA CTGGGAGGAG ACGGCGTCGT TTAGGTAAAA GACGAAGAAG CAGACGCAGA 660
 GAAGATCCGA ATTACCCGAA ACCGAACCGG AGCGGTTACA ATTTCTTCTT TGCTGAGAAA 720
 CATTGCAAGC TCAAATCACT TTATCCCAAC AAGGAGAGAG AGTTTACGAA ACTTATCGGA 780
 GAATCGTGGA GCAATCTCTC TACCGAAGAA CGAATGGTAA CAAATTATCT TTAAACCGT 840
 TACCGATTTA GTGATGAAAT TAGATTTGTA GTAAAT 876

(2) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1357 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

GGATCCCCGG GTACCATCAT AATACTCATC CTCCTTCTCA AGGTTGCTAC GTATTATCAA 60
 5 TATCTAGTAT ATACTTGTCT TTGTTCTATG CTTTATATCA TCATTTTATG ACAAAAAATG 120
 ATTAAGGTCT TAGTTAATGA TTATGTATAT GTGAAACTTA TATTTAGGGG CACAGTTTAA 180
 TTTCTGATGA TAATTGTCTA GTTAGCTTTA TGTACTTATC ATAAAAACCT TAGTGTTTAT 240
 10 CGCAATACTT TTCAAATATA GTGTAGAATC ATAATGGTCC CACTGTCATT ATGTTTGATG 300
 CAAATCTATT TGGATTTTGT TGGATAATAA ACCGATGACG TGGACCAGAC CAGTAGCTAT 360
 15 AAGATTTGGT TCACATAGAA ATTTTTTATA AGATAATGTA TCTAGGTTTG CTTATGATTA 420
 TACATGTGAT ATTTAATACA TGGCACAGGT TCGTCGAGTT TCACAGCCAT AGGTACAATA 480
 GAAGGCAAAT TCGATTGTGG TTATCTGGTA AAAGTTAAGT TGGGCTCAGA GATTCTTAAC 540
 20 GGCCTTCTTT ATCATTCGGC CCAGCCCGGC CCATCATCAT CTCCAACCGC TGTTCATAAC 600
 AATGCCGTTG TACCTTATGT TGAAACTGGG AGGAGACGGC GTCGTTTAGG TAAAAGACGA 660
 25 AGAAGCAGAC GCAGAGAAGA TCCGAATTAC CCGAAACCGA ACCGGAGCGG TTACAATTTC 720
 TTCTTTGCTG AGAAACATTG CAAGCTCAAA TCACTTTATC CCAACAAGGA GAGAGAGTTT 780
 ACGAAACTTA TCGGAGAATC GTGGAGCAAT CTCTCTACCG AAGAACGAAT GGTAACAAAT 840
 30 TATCTTTTAA ACCGTTACCG ATTTAGTGAT GAAATTAGAT TTGTATTGCA GCGATTCAT 900
 TTTAGATTCT CAAAAATATT CTCAGATGTG TGGGATTGA GTAGAGTTTA TGTGCGTTG 960
 35 GCATGATTTG AATAGTATGC AAGATTTTGT AGATTTTGCA TTCGTTTCATG TGTGTATGTG 1020
 TGATTGTAGC TTGATATGAT TTAACCTGTT AGTTAAATGT GCATAGACAA TAAGTAACAT 1080
 40 ACGAAGCGAG TCACTAAGCA TAAGAGTCAA CTTGTTTTGC TGAAAAGATA TCACTTATGA 1140
 TTTTCCAATC ATTTTAGCTT TTTGTCACT TGAGCTTAAT GATTCTTCTG AAATTCGATT 1200
 CTTTGTGTTG TTTATGTCAC ATTCTTTAGA ATTGAGAATC TAAGAAATGC TTACAGGATA 1260
 45 TGGTGAAACT ATTCTTTTAA GATAGCATGA TGCTTCTTTT ATGATTCTAC AGTGGCTAAG 1320
 TCATTTTTTT TTTGTTCTAT TCTTTGTAGC ACCATGG 1357

(2) INFORMATION FOR SEQ ID NO: 19:

50

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 34 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 55 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

65 GTCTTGATACA AATCTAATTT CATCACTAAA TCGG 34

(2) INFORMATION FOR SEQ ID NO: 20:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 34 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

TTAGATTGT ACAAGACCCT TCCTCTATAT AAGG

34

(2) INFORMATION FOR SEQ ID NO: 21:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

TTCCCACTCA CGACGTTGT

19

(2) INFORMATION FOR SEQ ID NO: 22:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1006 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

55	GGATCCCCGG GTACCATCAT AATACTCATC CTCCTTCTCA AGGTTGCTAC GTATTATCAA	60
	TATCTAGTAT AACTTGTCT TTGTCTATG CTTTATATCA TCATTTTATG ACAAAAAATG	120
	ATTAAGGTCT TAGTTAATGA TTATGTATAT GTGAACTTA TATTTAGGGG CACAGTTTAA	180
60	TTTCGTATGA TAATTGTCTA GTTAGCTTTA TGTACTTATC ATAAAAACCT TAGTGTTTAT	240
	CGCAATACTT TTCAAATATA GTGTAGAATC ATAATGGTCC CACTGTCATT ATGTTTGATG	300
65	CAAATCTATT TGGATTTTGT TGGATAATAA ACCGATGACG TGGACCAGAC CAGTAGCTAT	360

	AAGATTTGGT TCACATAGAA ATTTTATA AGATAATGTA TCTAGGTTTG CTTATGATT	420
	TACATGTGAT ATTTAATACA TGGCACAGGT TCGTCGAGTT TCACAGCCAT AGGTACAATA	480
5	GAAGGCAAAT TCGATTGTGG TTATCTGGTA AAAGTTAAGT TGGGCTCAGA GATTCTTAAC	540
	GGCGTTCTTT ATCATTCGGC CCAGCCCGGC CCATCATCAT CTCCAACCGC TGTTCTAAAC	600
10	AATGCCGTTG TACCTTATGT TGAACTGGG AGGAGACGGC GTCGTTTAGG TAAAAGACGA	660
	AGAAGCAGAC GCAGAGAAGA TCCGAATTAC CCGAAACCGA ACCGGAGCGG TTACAATTTC	720
	TTCTTTGCTG AGAAACATTG CAAGCTCAAA TCACTTTATC CCAACAAGGA GAGAGAGTTT	780
15	ACGAAACTTA TCGGAGAATC GTGGAGCAAT CTCTCTACCG AAGAACGAAT GGTAACAAAT	840
	TATCTTTTAA ACCGTTACCG ATTTAGTGAT GAAATTAGAT TTGTACAAGA CCCTTCCTCT	900
20	ATATAAGGAA GTTCATTTC TTTGGAGAGG ACACGTATTT TTACAACAAT TACCAACAAC	960
	AACAAACAAC AAACAACATT ACAATTACTA TTTACAATTA CCATGG	1006